

A21197

Leader in Biomolecular Solutions for Life Science



CD141/Thrombomodulin Rabbit mAb

Catalog No.: A21197 **Recombinant**

Basic Information

Observed MW

90kDa

Calculated MW

60kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

WB,IF/ICC,FC,ELISA

Cross-Reactivity

Human

CloneNo number

ARC52599

Recommended Dilutions

WB	1:1000 - 1:5000
IF/ICC	1:50 - 1:200
FC	1:50 - 1:200
ELISA	Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

Contact

 www.abclonal.com

Background

The protein encoded by this intronless gene is an endothelial-specific type I membrane receptor that binds thrombin. This binding results in the activation of protein C, which degrades clotting factors Va and VIIIa and reduces the amount of thrombin generated. Mutations in this gene are a cause of thromboembolic disease, also known as inherited thrombophilia.

Immunogen Information

Gene ID	Swiss Prot
7056	P07204

Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 19-515 of human CD141/Thrombomodulin (NP_000352.1).

Synonyms

TM; THRM; AHUS6; BDCA3; CD141; BDCA-3; THPH12; CD141/Thrombomodulin

Product Information


Source	Isotype	Purification
Rabbit	IgG	Affinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300,0.05% BSA,50% glycerol,pH7.3.

Validation Data



Western blot analysis of lysates from A-431 cells, using CD141/Thrombomodulin Rabbit mAb (A21197) at 1:2000 dilution.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Enhanced Kit (RM00021).


Exposure time: 90s.



Immunofluorescence analysis of A431 cells using CD141/Thrombomodulin Rabbit mAb (A21197) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of THP-1 cells using CD141/Thrombomodulin Rabbit mAb (A21197) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (AS007) at 1:500 dilution. Blue: DAPI for nuclear staining.



Flow cytometry: 1×10^6 Jurkat cells (negative control, left) and THP-1 cells (right) were surface-stained with CD141/Thrombomodulin Rabbit mAb (A21197, 10 µg/mL, green line) or Rabbit IgG isotype control (AC042, 10 µg/mL, blue line), followed by Alexa Fluor 647 conjugated goat anti-rabbit pAb (1:600 dilution) staining. Non-fluorescently stained cells were used as blank control (red line).