



NeoStain Poly DS Kit - for
Mouse and Rabbit
antibody on Human
tissue (BCIP/AEC)

NB-23-00087-1

NB-23-00087-2

NB-23-00087-3

NeoStain Poly DS Kit - for Mouse and Rabbit antibody on Human tissue (BCIP/AEC)

#Cat: NB-23-00087-1	Size: 12 mL(120 Slides)
#Cat: NB-23-00087-2	Size: 36 mL(360 Slides)
#Cat: NB-23-00087-3	Size: 120 mL(1200 Slides)

Storage: 4-8°C

Intended Use:

The **NeoStain Poly DS Kit** is designed to use with user supplied mouse and rabbit antibodies to detect two distinct antigens on human tissue or cell samples. This kit has been tested in paraffin tissue. However, this kit can be used on frozen specimen and freshly prepared monolayer cell smears.

Double staining is one of the most common methods used in immunohistochemistry to screen two distinct antigens in a single tissue ^{1, 2}. NeoBiotech Labs **NeoStain Poly DS Kit** supplies user with two polymer enzyme conjugates; an HRP-Polymer anti-Mouse IgG and AP-Polymer anti-Rabbit IgG with reactive chromogens for each enzyme. The AEC chromogen (Red Brick color) is used with HRP-Polymer anti-Mouse IgG and BCIP/NBT (Purple/Blue color) is used with AP-Polymer anti-Rabbit IgG. Simplified steps offer a much faster protocol as the enzyme conjugates are applied to the specimen as a mixture. Both the enzyme conjugated polymers and chromogens are optimized to give the strongest signal with no background. **NeoStain Poly DS Kit** is non-biotin system that avoids the need to block endogenous biotin causing non-specific binding.

Kit Components:

Component No.	Content	12mL Kit	36mL Kit	120mL Kit
Reagent 1	HRP-Polymer(AEC) anti-Mouse IgG (RTU)	6mL	18mL	60mL
Reagent 2	AP-Polymer anti-Rabbit IgG (RTU)	6mL	18mL	60mL
Reagent 3	BCIP/NBT (RTU)	12mL	18mLx2	120mL
Reagent 4A	AEC Substrate (20x)	1mL	2mL	6mL
Reagent 4B	AEC Chromogen (20x)	2mL	4mL	12mL
Reagent 4C	Hydrogen Peroxide (20x)	1mL	2mL	6mL
Reagent 5	Simpo-Mount (RTU)	12mL	18mLx2	120mL

Recommended Protocol:

1. Fixation: To ensure the quality of the staining and obtain reproducible performance, user needs to supply appropriately fixed tissue and well-prepared slides.
2. Tissues need to be adhered to the slide tightly to avoid tissue falling off.
3. Paraffin embedded section must be deparaffinized with xylene and rehydrated with a graded series of ethanol before staining.
4. Cell smear samples should be made as much monolayer as possible to obtain satisfactory results.
5. Three control slides will aid the interpretation of the result: positive tissue control, reagent control (slides treated with Isotype control reagent), and negative control.
6. Proceed IHC staining: DO NOT let specimen or tissue dry from this point on.
7. We recommend TBS-T to be used as the wash buffer to get the highest sensitivity and clean background. Phosphate in the PBS-T may inhibit the activity of the alkaline phosphatase.

Note: 1X TBS-T =50mM Tris HCl, 150mM NaCl, 0.05% Tween-20 pH7.6. NeoBiotech sells 10xTBS-T for your convenience **NB-23-00201**.

Reagent	Staining Procedure	Incubation Time(Min.)
1. Peroxidase and Alkaline Phosphatase Blocking Reagent Not provided We recommend using Dual Enzyme Block NB-23-00193-1 . Fast, easy and it will block endogenous alkaline phosphatase	a. Incubate slides in peroxidase and alkaline phosphatase blocking reagent. We recommend Dual Enzyme Block NB-23-00193-1 . b. Rinse the slide using distilled water.	10 min
2. HIER Pretreatment: Refer to antibody data sheet.	a. Heat Induced Epitope Retrieval (HIER) may be required for primary antibody suggested by vendor. b. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T (See note 7 above) ; 3 times for 2 minutes each.	
3. Preblock (optional)	For paraffin section, Improved formula saves the need for a preblock step. For frozen tissue, preblock may or may not be required depending on fixative.	
4. Mouse antibody 1 and Rabbit antibody 2: Supplied by user	Notes: Investigator needs to optimize dilution and incubation times prior to double staining. a. Apply 2 drops or enough volume of both Primary Antibody 1 and Antibody 2 to cover the tissue completely. Mix well on the slide and Incubate in moist chamber for 30-60 min. b. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T ; 3 times for 2 minutes each.	30-60 min
5. Reagent 1 and 2: Reagent 1: HRP Polymer(AEC) anti-Mouse IgG (RTU) Reagent 2: AP Polymer anti-Rabbit IgG (RTU)	a. Apply 1 drop (50µL) of Reagent 1 HRP Polymer(AEC) anti-Mouse IgG and 1 drop of Reagent 2 AP Polymer anti-Rabbit IgG to cover each section, mix well on the slide. Or you may prepare secondary antibodies cocktail in advance: 50µL Reagent 1 HRP Polymer(AEC) anti-Mouse IgG plus 50µL Reagent 2 AP Polymer anti-Rabbit IgG. b. Incubate in moist chamber for 30 min. c. Wash with 1X TBS-T only ; 3 times for 2 minutes each.	30 min
6. Reagent 3: BCIP/NBT (RTU)	a. Apply 2 drops or enough volume of Reagent 3 (BCIP/NBT) to completely cover tissue. Incubate for 3-10 min. b. Rinse thoroughly with distilled water. c. Wash with PBS-T containing 0.05% Tween-20 or 1X TBS-T ; 3 times for 2 minutes each.	5-10 min

<p>7. Reagent 4A, 4B, 4C: Reagent 4A:AEC Substrate (20x) Reagent 4B:AEC Chromogen (20x) Reagent 4C: Hydrogen Peroxide (20x)</p>	<p>a. Add 1 drop (50µL) of Reagent 4A to 1mL distilled water. Mix well . Add 2 drops of Reagent 4B and 1 drop of Reagent 4C to diluted reagent 1. Mix well. Keep away from light and use within 1 hour.</p> <p>b. Apply 2 drops (100µL) or enough volume of pre-mixed AEC solution to completely cover the tissue. Incubate for 5-15min, observe appropriate color development.</p> <p>c. Rinse well with distilled water. (AEC is alcohol soluble; do not dehydrate.)</p>	<p>10 min</p>
<p>8. HEMATOXYLIN Not provided</p>	<p>a. Counterstain with 2 drops (100µL) or enough volume of hematoxylin to completely cover tissue. Incubate for 10-15 seconds.</p> <p>b. Rinse thoroughly with tap water for 2-3 min.</p> <p>c. Put slides in PBS until show blue color (about ½ - 1 min.)</p> <p>d. Rinse well in distilled water.</p>	
<p>9. Reagent 5: Simpo-Mount(RTU)</p>	<p>a. Apply 2 drops (100µL) or enough volume Reagent 5 to cover tissue when tissue is wet. Rotate the slides to allow Simpo-Mount spread evenly. DO NOT coverslip.</p> <p>b. Place slides horizontally in an oven at 40-50°C for at least 30 minutes or leave it at room temperature until slides are thoroughly dried. Hardened Simpo-Mount forms an impervious polymer barrier to organic solvent. Do not use oil directly on the top of dried Simpo-Mount.</p>	<p>30 min in 40-50°C oven Or: overnight at room temperature</p>

Protocol Notes:

1. The fixation, tissue slide thickness, antigen retrieval and primary antibody dilution and incubation time affect results significantly. Investigator needs to consider all factors and determine optimal conditions when interpreting the result.
2. Simpo-Mount is an aqueous-based mounting media for immunohistochemistry. It is used as the permanent mounting media for alcohol soluble chromogens such as AP-Red, AEC, and BCIP. Simpo-Mount does not use a coverslip. However, if you need to coverslip your tissue, after Simpo-Mount has dried, dip the slide in xylene (1 to 2 seconds), apply an organic mounting solution (such as **NeoBio Mount Perm**, Cat# NB-23-00156), and place cover glass on the slide. Store slides after they have dried completely.

Precautions:

Please wear gloves and take other necessary precautions.

Remarks:

For research use only.

References:

1-[De Pasquale A](#), [Paterlini P](#), [Quaglino D](#). *Immunochemical demonstration of different antigens in single cells in paraffin-embedded histological sections*. [Clin Lab Haematol](#). 1982;4(3):267-72.

2-Polak J. M and Van Noorden S. *Introduction to Immunocytochemistry Second Edition*. Bios Scientific Publishers. P41-54. 1997