

**SHANK (pan) Antibody**  
**SHANK Antibody, Clone S23b-49**  
**Catalog # ASM10203****Specification**

---

**SHANK (pan) Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O9OX74</a>
Host	<b>Mouse</b>
Isotype	<b>IgG1</b>
Reactivity	<b>Human, Mouse, Rat</b>
Clonality	<b>Monoclonal</b>

**Description**

Mouse Anti-Rat SHANK (pan) Monoclonal IgG1

**Target/Specificity**

Detects ~160kDa. Recognizes Shank1, 2 and 3.

**Other Names**

Cortactin binding protein 1 Antibody, Cortactin SH3 domain-binding protein Antibody, Cortactin-binding protein 1 Antibody, CortBP1 Antibody, CTTNBP1 Antibody, GKAP/SAPAP interacting protein Antibody, GKAP/SAPAP-interacting protein Antibody, KIAA1022 Antibody, Proline rich synapse associated protein 1 Antibody, Proline-rich synapse-associated protein 1 Antibody, ProSAP1 Antibody, SH3 and multiple ankyrin repeat domains protein 2 Antibody, SHAN2\_RAT Antibody, SHANK Antibody, Shank2 Antibody, SPANK-3 Antibody

**Immunogen**

Fusion protein amino acids 84-309 of rat Shank2

**Purification**

Protein G Purified

Storage **-20°C****Storage Buffer**

PBS pH7.4, 50% glycerol, 0.09% sodium azide

Shipping Temperature **Blue Ice or 4°C****Certificate of Analysis**

1 µg/ml of SMC-327 was sufficient for detection of Shank1-4 in 10 µg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

**Cellular Localization**


Cytoplasm | Cell Junction | Synapse | Postsynaptic Cell Membrane | Postsynaptic Density | Cell Projection | Growth Cone

**SHANK (pan) Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**SHANK (pan) Antibody - Images**



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-SHANK (pan) Monoclonal Antibody, Clone S23b-49 (ASM10203). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-SHANK (pan) Monoclonal Antibody (ASM10203) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:200 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Cytoplasm . Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) SHANK (pan) Antibody (D) Composite.



Western Blot analysis of Rat brain membrane lysate showing detection of SHANK protein using

Mouse Anti-SHANK Monoclonal Antibody, Clone S23b-49 (ASM10203). Load: 15 µg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-SHANK Monoclonal Antibody (ASM10203) at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.

### **SHANK (pan) Antibody - Background**

Shank proteins make up a family of scaffold proteins identified through their interaction with a variety of membrane and cytoplasmic proteins (1). Shank proteins at postsynaptic sites of excitatory synapses play roles in signal transmission into the postsynaptic neuron. Studies suggest that Shank2 is expressed in the neurons of the developing retina, and could play a role in the neuronal differentiation of the developing retina (2). Other recent studies suggest that the disruption of glutamate receptors at the Shank postsynaptic platform could contribute to the destruction of the postsynaptic density, which underlies the synaptic dysfunction and loss in Alzheimer's disease (3).

### **SHANK (pan) Antibody - References**

1. Sheng M., and Kim E. (2000) *Journal of Cell Science*. 113: 1851-1856.
2. Kim J.H., et al. (2009) *Exp Mol Med*. 41(4): 236-242.
3. Gong Y., et al. (2009) *Brain Res*. 1292: 191-198.