

HSP60 Antibody

Catalog # ASM10360

Specification

HSP60 Antibody - Product Information

Application WB, IHC, IP
Primary Accession P10809
Other Accession NP_002147.2
Host Rabbit

Reactivity Human, Mouse, Rat, Rabbit, Chicken,

Bovine, Dog Polyclonal

Clonality

Description

Rabbit Anti-Human HSP60 Polyclonal

Target/Specificity
Detects ~60kDa.

Other Names

CPN60 Antibody, GROEL Antibody, HLD4 Antibody, HSP 60 Antibody, HSP65 Antibody, HSPD1 Antibody, HuCHA60 Antibody, SPG 13 Antibody

Immunogen

Human HSP60 produced through recombinant DNA methods in E.coli

PurificationProtein A Purified

Storage -20°C

Storage Buffer

PBS, 50% glycerol, 0.09% sodium azide

Shipping Temperature Blue Ice or 4°C

Certificate of Analysis

 $1 \mu g/ml$ of SPC-105 was sufficient for detection of HSP60 in 20 μg of heat shocked HeLa cell lysate by colorimetric immunoblot analysis using goat anti-mouse IgG as the secondary antibody.

Cellular Localization

Mitochondrion | Mitochondrion Matrix

HSP60 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence



Tel: 858.875.1900 Fax: 858.875.1999

- Immunoprecipitation
- Flow Cytomety
- Cell Culture

HSP60 Antibody - Images

HSP60 Antibody - Background

In both prokaryotic and eukaryotic cells, the misfolding and aggregation of proteins during biogenesis and under conditions of cellular stress are prevented by molecular chaperones. Members of the HSP60 family of heat shock proteins are some of the best characterized chaperones. HSP60, also known as Cpn60 or GroEl, is an abundant protein synthesized constitutively in the cell that is induced to a higher concentration after brief cell shock. It is present in many species and exhibits a remarkable sequence homology among various counterparts in bacteria, plants, and mammals with more than half of the residues identical between bacterial and mammalian HSP60 (1-3). Whereas mammalian HSP60 is localized within the mitochondria, plant HSP60, or otherwise known as Rubisco-binding protein, is located in plant chloroplasts. It has been indicated that these proteins carry out a very important biological function due to the fact that HSP60 is present in so many different species. The common characteristics of the HSP60s from the divergent species are i) high abundance, ii) induction with environmental stress such as heat shock, iii) homo-oligomeric structures of either 7 or 14 subunits which reversibly dissociate in the presence of Mg2+ and ATP, iv) ATPase activity and v) a role in folding and assembly of oligomeric protein structures (4). These similarities are supported by recent studies where the single-ring human mitochondrial homolog, HSP60 with its co-chaperonin, HSP10 were expressed in a E. coli strain, engineered so that the groE operon is under strict regulatory control. This study has demonstrated that expression of HSP60-HSP10 was able to carry out all essential in vivo functions of GroEL and its co-chaperonin, GroES (5). HSP60 has however been linked to a number of autoimmune diseases, as well as Alzheimer's, coronary artery diseases, MS, and diabetes (6-9).

HSP60 Antibody - References

- 1. Hartl, F.U. (1996) Nature 381: 571-579.
- 2. Bukau, B. and Horwich, A.L. (1998) Cell 92: 351-366.
- 3. Hartl, F.U. and Hayer-Hartl, M. (2002) Science 295: 1852- 1858.
- 4. Jindal, S., et al. (1989) Molecular and Cellular Biology 9: 2279-2283.
- 5. La Verda, D., et al (1999) Infect Dis. Obstet. Gynecol. 7: 64-71.
- 6. Itoh, H. et al. (2002) Eur. J. Biochem. 269: 5931-5938.
- 7. Gupta, S. and Knowlton, A.A. J. Cell Mol Med. 9: 51-58.
- 8. Deocaris, C.C. et al. (2006) Cell Stress Chaperones 11: 116-128.
- 9. Lai, H.C. et al. (2007) Am. J. Physiol. Endocrinol. Metab. 292: E292-E297.
- 10. Gao, Y.L., et al (1995) J. of Immunology 154: 3548-3556.
- 11. Neuer, A., et al (1997) European Society for Human Reproduction and Embryology 12(5):925-929.
- 12. Bason, C., et al (2003) Lancet 362(9400): 1971-1977.