

mGluR1/5 Antibody

mGluR1/5 Antibody, Clone S75-33 Catalog # ASM10241

Specification

mGluR1/5 Antibody - Product Information

Application Primary Accession Other Accession

Host Isotype Reactivity Clonality

Format

Description

Mouse Anti-Rat mGluR1/5 Monoclonal IgG1

Mouse Anti-Nat Miglak1/3 Monociol

Target/Specificity

Detects ~130kDa (both mGluR1 and mGluR5).

Other Names

Glutamate receptor metabotropic 5, GPRC1E Antibody, GRM 5 Antibody, GRM5 Antibody, Metabotropic glutamate receptor 5 Antibody, Metabotropic glutamate receptor 5 variant F Antibody, Metabotropic glutamate receptor 5 variant G Antibody, Metabotropic glutamate receptor 5 variant H Antibody, mGlu5 Antibody, mGlu85 Antibody, mGlu85 Antibody, mGlu85 Antibody, Glutamate receptor metabotropic 1 Antibody, glutamate receptor Antibody, metabotropic 1 Antibody, GRM1 Antibody, GRM1 Antibody, GRM1 Antibody, GRM1 Antibody, mGlu81 Antibody, mGlu81 Antibody, mGlu81 Antibody, mGlu81 Antibody, mGlu81 Antibody, mGlu81 Antibody

WB

P31424

Mouse

IqG1

FITC

NP 058708.1

Monoclonal

Human, Mouse, Rat

Immunogen

Fusion protein amino acids 824-1203 (cytoplasmic C-terminus) of rat mGluR5b

Purification

Protein G Purified

Storage -20°C

Storage Buffer

PBS pH7.4, 50% glycerol, 0.09% sodium azide

Shipping Temperature

Blue Ice or 4ºC

Certificate of Analysis

1 μ g/ml of SMC-407 was sufficient for detection of mGluR1/5 glutamate receptor in 20 μ g of rat brain membrane lysate and assayed by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.

Cellular Localization

Cell Membrane


mGluR1/5 Antibody - Protocols




Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

mGluR1/5 Antibody - Images




Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-mGluR1/5 Monoclonal Antibody, Clone N75/33 (ASM10241). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4% PFA for 15 min. Primary Antibody: Mouse Anti-mGluR1/5 Monoclonal Antibody (ASM10241) at 1:100 for overnight at 4°C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain; Hoechst (blue) nuclear stain at 1:800, 1.6mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) mGluR1/5 Antibody (D) Composite.





Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-mGluR1/5 Monoclonal Antibody, Clone N75/33 (ASM10241). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-mGluR1/5 Monoclonal Antibody (ASM10241) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:200 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Cell Membrane, Cytoplasm, Nucleus. Magnification: 60X. (A) DAPI (blue) nuclear stain. (B) Phalloidin Texas Red F-Actin stain. (C) mGluR1/5 Antibody. (D) Composite.



Western Blot analysis of Rat brain membrane lysate showing detection of mGluR5 Glutamate Receptor protein using Mouse Anti-mGluR5 Glutamate Receptor Monoclonal Antibody, Clone N75/33 (ASM10241). Primary Antibody: Mouse Anti-mGluR5 Glutamate Receptor Monoclonal Antibody (ASM10241) at 1:1000.

mGluR1/5 Antibody - Background

The AMPA receptor is a non-NMDA-type ionotropic transmembrane receptor for glutamate that mediates fast synaptic transmission in the CNS. AMPARs are composed of four types of subunits, designated as GluR1, GluR2, GluR3 and GluR4, which combine to form tetramers (1,2).

mGluR1/5 Antibody - References

- 1. Mayer M.L. (2005) Current Opinion in Neurobio. 15(3): 282-288.
- 2. Keinane K., et al. (1990) Science. 249: 556-560.