

**SHANK1/SHANK3 Antibody**  
**SHANK1/SHANK3 Antibody, Clone S367-51**  
**Catalog # ASM10292****Specification****SHANK1/SHANK3 Antibody - Product Information**

Application	<b>WB, IHC, ICC</b>
Primary Accession	<a href="#">O9JLU4</a>
Other Accession	<a href="#">NP_067708.1</a>
Host	<b>Mouse</b>
Isotype	<b>IgG2a</b>
Reactivity	<b>Human, Mouse, Rat</b>
Clonality	<b>Monoclonal</b>

**Description**

Mouse Anti-Rat SHANK1/SHANK3 Monoclonal IgG2a

**Target/Specificity**

Detects ~190kDa. Cross-reacts with SHANK1. Does not cross-react with SHANK2.

**Other Names**

Shank postsynaptic density protein Antibody, SH3 and multiple ankyrin repeat domains 3 Antibody, Proline rich synapse associated protein 2 Antibody, Proline-rich synapse-associated protein 2 Antibody, ProSAP2 Antibody, PSAP2 Antibody, Antibody, SH3 and multiple ankyrin repeat domains protein 3 Antibody, SH3/ankyrin domain gene 3, Shank3b Antibody, SPANK 2 Antibody, SPANK2 Antibody, KAP/SAPAP interacting protein Antibody, OTTHUMP00000174437 Antibody, SH3 and multiple ankyrin repeat domains 1 Antibody, SH3 and multiple ankyrin repeat domains protein 1 Antibody, SH3/ankyrin domain gene 1 Antibody, SHANK 1 Antibody, Shank1 Antibody, Shank1a Antibody, Somatostatin receptor interacting protein Antibody, Somatostatin receptor-interacting protein Antibody, SPANK 1 Antibody, SPANK1 Antibody, SSTR interacting protein Antibody, SSTR-interacting protein Antibody, SSTRIP Antibody, Synamon Antibody, A1841104 Antibody, DEL22q13.3 Antibody, KIAA1650 Antibody

**Immunogen**

Fusion protein amino acids 538-626 (SH3 domain) of rat SHANK3. Mouse: 100% identity (89/89 amino acids identical). Human: 97% identity (87/89 amino acids identical). ~70% identity with SHANK1 and SHANK2.

**Purification**

Protein G Purified

Storage **-20°C**

**Storage Buffer**

PBS pH 7.4, 50% glycerol, 0.1% sodium azide

Shipping Temperature **Blue Ice or 4°C**

**Certificate of Analysis**

1 µg/ml of SMC-460 was sufficient for detection of SHANK1/SHANK3 in 20 µg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

**Cellular Localization**


Cytoplasm | Cell Junction

## SHANK1/SHANK3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## SHANK1/SHANK3 Antibody - Images



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-SHANK1/SHANK3 Monoclonal Antibody, Clone N367/51 (ASM10292). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4% PFA for 15 min. Primary Antibody: Mouse Anti-SHANK1/SHANK3 Monoclonal Antibody (ASM10292) at 1:100 for overnight at 4°C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain; Hoechst (blue) nuclear stain at 1:800, 1.6mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) SHANK1/SHANK3 Antibody (D) Composite.

## SHANK1/SHANK3 Antibody - Background

SHANK proteins are scaffolding adaptors that have been shown to integrate neurotransmitter receptors into the cortical cytoskeleton at postsynaptic densities. SHANK1-3 of the SHANK/ProSAP family are molecular scaffolds in the postsynaptic density (PSD). SHANK recruits betaPIX and PAK to dendritic spines to regulate postsynaptic structure and interacts with ionotropic receptor and metabotropic glutamate receptor complexes. Transcript splice variation in the Shank family influences the spectrum of Shank-interacting proteins in the PSDs of adult and developing brain to ensure normal development.