

VEGFR-1	Human, Recombinant, <i>E. coli</i>		
Cat. No.: NB-02-11 NB-02-11			
Synonym:	Vascular endothelial growth factor receptor-1 [precursor]; VEGFR-1; EC 2.7.1.112; Vascular permeability factor receptor ; Tyrosine-protein kinase receptor FLT; FLT-1; Tyrosine-protein kinase FRT; FMS-like tyrosine kinase 1.		
Description:	VEGFR-1 belongs to a family of five receptor tyrosine kinase receptors that are specific to a number of angiogenic factors including VEGF and PDGF, and are therefore thought to play a role in the angiogenic process. VEGFR- 1 is expressed in two forms, predominantly on endothelial cells, giving rise to a full-length membrane bound receptor and a truncated soluble receptor. Elevated levels of the soluble form of this VEGF receptor may be used in the diagnosis of pathological conditions.		
	Recombinant VEGFR-1 comprises a 298 amino acid fragment (31-328) corresponding to the IgG-like domains 1-3 from the mature soluble VEGFR protein. VEGF-R1 is expressed in <i>E. coli</i> with an amino-terminal hexahistidine tag. This product is for research use only and is not intended for diagnostic or therapeutic use.		
Form:	Liquid	148 kDa	
Purity:	>95% by SDS-PAGE	64 kDa 50 kDa ← VEGFR-1	
		36 kDa	
		22 kDa	
		16 kDa	
Reference(s):	Starovasnik, M. A <i>et al</i> . (1999) <i>J. Mol. Biol.</i> 296 (3) : 531-544.		

Kanno, S. et al. (2000) Oncogene 19 (17): 2138-2146.