# ABclonal www.abclonal.com

## NFIB / NF1B2 Rabbit mAb

Catalog No.: A9125 Recombinant

## **Basic Information**

#### **Observed MW**

47kDa

#### **Calculated MW**

47kDa

#### **Category**

SMab Recombinant Monoclonal Antibody

### **Applications**

WB, ELISA

#### **Cross-Reactivity**

Human, Mouse

#### CloneNo number

ARC1859

## **Background**

Enables DNA-binding transcription activator activity, RNA polymerase II-specific; RNA polymerase II cis-regulatory region sequence-specific DNA binding activity; and transcription regulator inhibitor activity. Involved in brain development; negative regulation of DNA binding activity; and regulation of transcription by RNA polymerase II. Located in fibrillar center and nucleoplasm.

## **Recommended Dilutions**

**WB** 

1:500 - 1:1000

## **Immunogen Information**

Gene ID 4781 Swiss Prot

## Immunogen

A synthetic peptide corresponding to a sequence within amino acids 321-420 of human NFIB / NF1B2 (NP\_005587.2).

#### **Synonyms**

CTF; MACID; NF1-B; NFI-B; NFIB2; NFIB3; NF-I/B; NFI-RED; HMGIC/NFIB; NFIB / NF1B2

#### **Contact**

•

www.abclonal.com

## **Product Information**


SourceIsotypePurificationRabbitIgGAffinity purification

## Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.02% sodium azide, 0.05% BSA, 50% glycerol, pH7.3.

## **Validation Data**



Western blot analysis of extracts of various cell lines, using NFIB / NF1B2 antibody (A9125) at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 180s.