

GRP78 Antibody
GRP78 Antibody, Clone 3C5-1A4
Catalog # ASM10155

Specification

GRP78 Antibody - Product Information

Application	WB
Primary Accession	P06761
Other Accession	NP_037215.1
Host	Mouse
Isotype	IgG1 Kappa
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal

Description

Mouse Anti-Rat GRP78 Monoclonal IgG1 Kappa

Target/Specificity

Detects ~78kDa.

Other Names

78 kDa glucose regulated protein Antibody, 78 kDa glucose-regulated protein Antibody, AL022860 Antibody, AU019543 Antibody, BIP Antibody, D2Wsu141e Antibody, D2Wsu17e Antibody, Endoplasmic reticulum luminal Ca(2+)-binding protein grp78 Antibody, Endoplasmic reticulum luminal Antibody, Ca²⁺ binding protein grp78 Antibody, FLJ26106 Antibody, Glucose Regulated Protein 78kDa Antibody, GRP 78 Antibody, GRP-78 Antibody, GRP78_HUMAN Antibody, Heat shock 70 kDa protein 5 Antibody, Heat Shock 70kDa Protein 5 Antibody, H5Ce70 Antibody, HSPA 5 Antibody, HSPA5 Antibody, Immunoglobulin Heavy Chain Binding Protein Antibody, Immunoglobulin heavy chain-binding protein Antibody, mBiP Antibody, MIF2 Antibody, Sez7 Antibody

Immunogen

Full-length recombinant rat GRP78

Purification

Protein G Purified

Storage **-20°C**

Storage Buffer

PBS pH7.4, 50% glycerol, 0.09% sodium azide

Shipping Temperature **Blue Ice or 4°C**

Certificate of Analysis

1 µg/ml of SMC-210 was sufficient for detection of GRP78 in 20 µg of HEK-293 lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

Cellular Localization


Endoplasmic Reticulum | Endoplasmic Reticulum Lumen | Melanosome

GRP78 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GRP78 Antibody - Images



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-GRP78 Monoclonal Antibody, Clone 3C5-1A4 (ASM10155). Tissue: Fibroblast cell line (NIH 3T3). Species: Mouse. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-GRP78 Monoclonal Antibody (ASM10155) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: DAPI (blue) nuclear stain at 1:5000 for 5 min RT. Localization: Nucleus. Magnification: 60X.



Western Blot analysis of Human, Mouse, Rat NIH3T3, Rat Brain, and HEK-293 cell lysates showing detection of ~78 kDa GRP78 protein using Mouse Anti-GRP78 Monoclonal Antibody, Clone 3C5-1A4 (ASM10155). Lane 1: MW ladder. Lane 2: Mouse NIH3T3. Lane 3: Rat Brain. Lane 4: Human HEK-293. Block: 5% milk + TBST for 1 hour at RT. Primary Antibody: Mouse Anti-GRP78 Monoclonal Antibody (ASM10155) at 1:1000 for 1 hour at RT. Secondary Antibody: HRP Goat Anti-Mouse at 1:50 for 1 hour at RT. Color Development: TMB solution for 5 min at RT. Predicted/Observed Size: ~78 kDa.

GRP78 Antibody - Background

GRP78 is a ubiquitously expressed, 78-kDa glucose-regulated protein, and is commonly referred to as an immunoglobulin chain binding protein (BiP). The BiP proteins are categorized as stress response proteins because they play an important role in the proper folding and assembly of nascent protein

and in the scavenging of misfolded proteins in the endoplasmic reticulum lumen. Translation of BiP is directed by an internal ribosomal entry site (IRES) in the 5' nontranslated region of the BiP mRNA. BiP IRES activity increases when cells are heat stressed (1). GRP78 is also critical for maintenance of cell homeostasis and the prevention of apoptosis (2). Luo et al. have provided findings that suggest GRP78 is essential for embryonic cell growth and pluripotent cell survival (3). In terms of diseases, GRP78 has been shown to be a reliable biomarker of hypoglycemia, to serve a neuroprotective function in neurons exposed to glutamate and oxidative stress (4), and its protein levels are reduced in the brains of Alzheimer's patients (5). Also, the induction of the GRP78 protein that results in severe glucose and oxygen deprivation could possibly lead to drug resistance to anti-tumor drugs (6, 7).

GRP78 Antibody - References

1. Cho, S. et al. (2007). *Mol Cell Biol* 27(1): 368-83.
2. Yang, Y. et al. (1998) *J Biol Chem* 273: 25552-25555.
3. Luo, S. et al (2006) 26 (15): 5688-97.
4. Yu, Z. et al. (1999) *Exp Neurol*. 15: 302-314.
5. Koomagi, R. et al. (1999) *Anticancer Res*. 19: 4333-4336.
6. Laquerre, S. et al. (1998) *J. Virology* 72: 4940-4949.
7. Dong, D. et al. (2005) *Cancer Res* 65(13): 5785-91.