

**GRP78 Antibody**  
**GRP78 Antibody, Clone 3C5-1A4**  
**Catalog # ASM10155**

**Specification**

---

**GRP78 Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">P06761</a>
Other Accession	<a href="#">NP_037215.1</a>
Host	<b>Mouse</b>
Isotype	<b>IgG1 Kappa</b>
Reactivity	<b>Human, Mouse, Rat</b>
Clonality	<b>Monoclonal</b>

**Description**

Mouse Anti-Rat GRP78 Monoclonal IgG1 Kappa

**Target/Specificity**

Detects ~78kDa.

**Other Names**

78 kDa glucose regulated protein Antibody, 78 kDa glucose-regulated protein Antibody, AL022860 Antibody, AU019543 Antibody, BIP Antibody, D2Wsu141e Antibody, D2Wsu17e Antibody, Endoplasmic reticulum luminal Ca(2+)-binding protein grp78 Antibody, Endoplasmic reticulum luminal Antibody, Ca2+ binding protein grp78 Antibody, FLJ26106 Antibody, Glucose Regulated Protein 78kDa Antibody, GRP 78 Antibody, GRP-78 Antibody, GRP78\_HUMAN Antibody, Heat shock 70 kDa protein 5 Antibody, Heat Shock 70kDa Protein 5 Antibody, H5Ce70 Antibody, HSPA 5 Antibody, HSPA5 Antibody, Immunoglobulin Heavy Chain Binding Protein Antibody, Immunoglobulin heavy chain-binding protein Antibody, mBiP Antibody, MIF2 Antibody, Sez7 Antibody

**Immunogen**

Full-length recombinant rat GRP78

**Purification**

Protein G Purified

Storage **-20°C**

**Storage Buffer**

PBS pH7.4, 50% glycerol, 0.09% sodium azide

Shipping Temperature **Blue Ice or 4°C**

**Certificate of Analysis**

1 µg/ml of SMC-210 was sufficient for detection of GRP78 in 20 µg of HEK-293 lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

**Cellular Localization**


Endoplasmic Reticulum | Endoplasmic Reticulum Lumen | Melanosome

**GRP78 Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### GRP78 Antibody - Images



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-GRP78 Monoclonal Antibody, Clone 3C5-1A4 (ASM10155). Tissue: Fibroblast cell line (NIH 3T3). Species: Mouse. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-GRP78 Monoclonal Antibody (ASM10155) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: DAPI (blue) nuclear stain at 1:5000 for 5 min RT. Localization: Nucleus. Magnification: 60X.



Western Blot analysis of Human, Mouse, Rat NIH3T3, Rat Brain, and HEK-293 cell lysates showing detection of ~78 kDa GRP78 protein using Mouse Anti-GRP78 Monoclonal Antibody, Clone 3C5-1A4 (ASM10155). Lane 1: MW ladder. Lane 2: Mouse NIH3T3. Lane 3: Rat Brain. Lane 4: Human HEK-293. Block: 5% milk + TBST for 1 hour at RT. Primary Antibody: Mouse Anti-GRP78 Monoclonal Antibody (ASM10155) at 1:1000 for 1 hour at RT. Secondary Antibody: HRP Goat Anti-Mouse at 1:50 for 1 hour at RT. Color Development: TMB solution for 5 min at RT. Predicted/Observed Size: ~78 kDa.

### GRP78 Antibody - Background

GRP78 is a ubiquitously expressed, 78-kDa glucose-regulated protein, and is commonly referred to as an immunoglobulin chain binding protein (BiP). The BiP proteins are categorized as stress response proteins because they play an important role in the proper folding and assembly of nascent protein

and in the scavenging of misfolded proteins in the endoplasmic reticulum lumen. Translation of BiP is directed by an internal ribosomal entry site (IRES) in the 5' nontranslated region of the BiP mRNA. BiP IRES activity increases when cells are heat stressed (1). GRP78 is also critical for maintenance of cell homeostasis and the prevention of apoptosis (2). Luo et al. have provided findings that suggest GRP78 is essential for embryonic cell growth and pluripotent cell survival (3). In terms of diseases, GRP78 has been shown to be a reliable biomarker of hypoglycemia, to serve a neuroprotective function in neurons exposed to glutamate and oxidative stress (4), and its protein levels are reduced in the brains of Alzheimer's patients (5). Also, the induction of the GRP78 protein that results in severe glucose and oxygen deprivation could possibly lead to drug resistance to anti-tumor drugs (6, 7).

### **GRP78 Antibody - References**

1. Cho, S. et al. (2007). *Mol Cell Biol* 27(1): 368-83.
2. Yang, Y. et al. (1998) *J Biol Chem* 273: 25552-25555.
3. Luo, S. et al (2006) 26 (15): 5688-97.
4. Yu, Z. et al. (1999) *Exp Neurol*. 15: 302-314.
5. Koomagi, R. et al. (1999) *Anticancer Res*. 19: 4333-4336.
6. Laquerre, S. et al. (1998) *J. Virology* 72: 4940-4949.
7. Dong, D. et al. (2005) *Cancer Res* 65(13): 5785-91.