

catalase siRNA (chicken): sc-270570

BACKGROUND

Catalase is a peroxisome specific marker protein belonging to the catalase family. Defects in the gene encoding for the catalase protein, CAT, can cause acatalasia, a disease characterized by the absence of catalase activity in red cells and associated with ulcerating oral lesions. Catalase is also an important regulator of oxidative stress and inflammation, and may contribute to the development of rheumatoid arthritis. Catalase, which can form a homotrimer, is found in nearly all aerobically respiring organisms and functions in protecting cells from the toxic effects of hydrogen peroxide.

REFERENCES

- Rodriguez-Esparragon, F.J., et al. 2003. Peroxisome proliferator-activated receptor- γ 2-Pro12Ala and endothelial nitric oxide synthase-4a/b gene polymorphisms are associated with essential hypertension. *J. Hypertens.* 21: 1649-1655.
- Rosmond, R., et al. 2003. The Pro12Ala PPAR γ 2 gene missense mutation is associated with obesity and Insulin resistance in Swedish middle-aged men. *Diabetes Metab. Res. Rev.* 19: 159-163.
- El-Sohemy, A., et al. 2005. Catalase and PPAR γ 2 genotype and risk of rheumatoid arthritis in Koreans. *Rheumatol. Int.* 26: 388-392.
- Eny, K.M., et al. 2005. Catalase and PPAR γ 2 genotype and risk of systemic lupus erythematosus in Koreans. *Lupus* 14: 351-355.
- Zhao, X., et al. 2006. 15d-Prostaglandin J₂ activates peroxisome proliferator-activated receptor- γ , promotes expression of catalase, and reduces inflammation, behavioral dysfunction, and neuronal loss after intracerebral hemorrhage. *J. Cereb. Blood Flow Metab.* 26: 811-820.
- SWISS-PROT/TrEMBL (P04040). World Wide Web URL: <http://www.expasy.ch/sprot/sprot-top.html>

CHROMOSOMAL LOCATION

Genetic locus: CAT (chicken) mapping to 5.

PRODUCT

catalase siRNA (chicken) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see catalase shRNA Plasmid (chicken): sc-270570-SH and catalase shRNA (chicken) Lentiviral Particles: sc-270570-V as alternate gene silencing products.

For independent verification of catalase (chicken) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-270570A, sc-270570B and sc-270570C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

catalase siRNA (chicken) is recommended for the inhibition of catalase expression in chicken cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

catalase (H-9): sc-271803 is recommended as a control antibody for monitoring of catalase gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor catalase gene expression knockdown using RT-PCR Primer: catalase (chicken)-PR: sc-270570-PR (20 μ l, 449 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.