PE Rabbit anti-Human Integrin β5/ITGB5 mAb

Catalog No.: A24597



Basic Information

Observed MW Refer to figures

Calculated MW 88kDa

Category

SMab Recombinant Monoclonal Antibody

Applications FC

Cross-Reactivity Human

CloneNo number ARC60858-PE

Conjugate

PE. Ex:565nm. Em:574nm.

Recommended Dilutions

FC

5 µl per 10^6 cells in 100 µl volume

Background

Integrins, heterodimeric trans-membrane matrix receptors, are mainly involved in the signal transduction and attachments to extra cellular matrix (ECM).⁺⁺ In ECM they act as a mediator of cell adhesion.⁺ In human, at least 18 α and eight β subunits have been found.⁺ ITGB5 (integrin subunit β 5) helps in facilitation of cancer cell migration, anchorage-independent growth and tumor angiogenesis. Integrin is activated by G-protein-coupled receptors, resulting in the phosphorylation of cytoplasmic domain of the β subunit. The coupled α and β cytoplasmic tails are responsible for maintaining integrin in inactive state.

Immunogen Information

Gene ID 3693

Swiss Prot P18084

Immunogen

Recombinant Protein corresponding to a sequence within amino acids 24-490 of human Integrin β 5/ITGB5 (NP_002204.2).

Synonyms

ITGB5; integrin beta-5

Product Information

Contact

€


Source Rabbit **Isotype** IgG **Purification** Affinity purification


www.abclonal.com

Storage

Store at 2-8°C. Avoid freeze. Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.

Validation Data





Flow cytometry:1X10^6 Jurkat cells (negative control,left) and A549 cells (right) were surface-stained with PE Rabbit anti-Human Integrin β 5/ITGB5 mAb(A24597,5 μ I/Test,orange line) or PE Rabbit IgG isotype control (5 μ I/Test,blue line). Non-fluorescently stained cells were used as blank control (red line). Flow cytometry:1X10^6 A549 cells cells were surface-stained with PE Rabbit IgG isotype control (5 μ l/Test,left) or PE Rabbit anti-Human Integrin β 5/ITGB5 mAb(A24597,5 μ l/Test,right).