ABclonal www.abclonal.com

PE Rabbit anti-Human/Monkey CD20 mAb

Catalog No.: A24420

Basic Information

Observed MW

Refer to figures

Calculated MW

14kDa/33kDa

Category

SMab Recombinant Monoclonal Antibody

Applications

FC

Cross-Reactivity

Human

CloneNo number

ARC51683-PE

Conjugate

PE. Ex:565nm. Em:574nm.

Recommended Dilutions

FC

5 μ l per 10^6 cells in 100 μ l volume

Background

This gene encodes a member of the membrane-spanning 4A gene family. Members of this nascent protein family are characterized by common structural features and similar intron/exon splice boundaries and display unique expression patterns among hematopoietic cells and nonlymphoid tissues. This gene encodes a B-lymphocyte surface molecule which plays a role in the development and differentiation of B-cells into plasma cells. This family member is localized to 11q12, among a cluster of family members. Alternative splicing of this gene results in two transcript variants which encode the same protein.

Immunogen Information

Gene ID931

Swiss Prot
P11836

Immunogen

A synthetic peptide corresponding to a sequence within amino acids 100-200 of human CD20 mAb (NP_068769.2).

Synonyms

MS4A1; B1; Bp35; CD20; CVID5; LEU-16; MS4A2; S7; B-lymphocyte antigen CD20

Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at 2-8°C. Avoid freeze.


Buffer: PBS with 0.03% proclin300,0.2% BSA,pH7.3.


Contact

€

<u>www.abclonal.com</u>

Validation Data





Flow cytometry:1X10^6 Jurkat cells (negative control,left) and Daudi cells (right) were surface-stained with PE Rabbit anti-Human/Monkey CD20 mAb(A24420,5 µl/Test,orange line) or PE Rabbit IgG isotype control (5 µl/Test,blue line). Non-fluorescently stained cells were used as blank control (red line).

Flow cytometry:1X10^6 Daudi cells were surface-stained with PE Rabbit IgG isotype control (5 μ l/Test,left) or PE Rabbit anti-Human/Monkey CD20 mAb(A24420,5 μ l/Test,right).