

AC035

Leader in Biomolecular Solutions for Life Science



# HRP-conjugated GAPDH Mouse mAb

Catalog No.: AC035

18 Publications

## Basic Information

### Observed MW

36kDa

### Calculated MW

36kDa

### Category

Mouse Monoclonal Antibody

### Applications

WB, ELISA

### Cross-Reactivity

Human, Mouse, Rat

### CloneNo number

AMC0062R

### Conjugate

HRP

## Recommended Dilutions

WB 1:2000 - 1:10000

## Background

This gene encodes a member of the glyceraldehyde-3-phosphate dehydrogenase protein family. The encoded protein has been identified as a moonlighting protein based on its ability to perform mechanistically distinct functions. The product of this gene catalyzes an important energy-yielding step in carbohydrate metabolism, the reversible oxidative phosphorylation of glyceraldehyde-3-phosphate in the presence of inorganic phosphate and nicotinamide adenine dinucleotide (NAD). The encoded protein has additionally been identified to have uracil DNA glycosylase activity in the nucleus. Also, this protein contains a peptide that has antimicrobial activity against *E. coli*, *P. aeruginosa*, and *C. albicans*. Studies of a similar protein in mouse have assigned a variety of additional functions including nitrosylation of nuclear proteins, the regulation of mRNA stability, and acting as a transferrin receptor on the cell surface of macrophage. Many pseudogenes similar to this locus are present in the human genome. Alternative splicing results in multiple transcript variants.

## Immunogen Information

### Gene ID

2597

### Swiss Prot

P04406

### Immunogen

Recombinant fusion protein containing a sequence corresponding to amino acids 1-335 of human GAPDH (NP\_002037.2).

### Synonyms

G3PD; GAPD; HEL-S-162eP; HRP-conjugated GAPDH

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Mouse

### Isotype

IgG1, Kappa

### Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.05% proclin300, 50% glycerol, pH7.3

## Validation Data



Western blot analysis of various lysates, using HRP-conjugated GAPDH Mouse mAb (AC035) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 10s.