


Octyl Glucose Neopentyl Glycol (OGNG) - Purity: =98% all anomers (by HPLC analysis) - = 4% anomers other than 1,3-bis-β (HPLC) [1257853-32-9]

#Cat: NB-19-0054-1g	Size: 1g
#Cat: NB-19-0054-5g	Size: 5g
#Cat: NB-19-0054-25g	Size: 25g



Description

Octyl glucose neopentyl glycol (OGNG) is a neopentyl glycol detergent that can be used to maintain the stability of membrane proteins[1].

In vitro, the decisive step, in the improvement of *Thermotoga maritima* (TmPPase) crystal quality came through detergent exchange in to Octyl glucose neopentyl glycol, a neopentyl glycol detergent[1].

Product Information

Codes	NB-19-0054-1G, NB-19-0054-5G, NB-19-0054-25G
Sizes	1g, 5g, 25g
Synonyms	2,2-dihexylpropane-1,3-bis-β-D-glucopyranoside, OGNG
CAS number	1257853-32-9
Formula	C ₂₇ H ₅₂ O ₁₂
Molecular Weight	568.69
Purity (HPLC)	Min 98%
Note	For research use only.

Description

In vitro :

Preparing stock solutions (Volume of Solvent) :

Mass	1 mg	5 mg	10 mg
Concentration			
1 mM	1.7584 mL	8.7921 mL	17.5843 mL
5 mM	0.3517 mL	1.7584 mL	3.5169 mL
10mM	0.1758 mL	0.8792 mL	1.7584 mL

DMSO : 100 mg/mL (175.84 mM; Need ultrasonic)

Please refer to the solubility information to select the appropriate solvent.

In vitro :

1. Add each solvent one by one: 10% DMSO >> 40% PEG300 >> 5% Tween-80 >> 45% saline
Solubility: ≥ 5 mg/mL (8.79 mM); Clear solution
2. Add each solvent one by one: 10% DMSO >> 90% (20% SBE- β -CD in saline) Solubility: ≥ 5 mg/mL (8.79 mM); Clear solution
3. Add each solvent one by one: 10% DMSO >> 90% corn oil
Solubility: ≥ 5 mg/mL (8.79 mM); Clear solution

References

- [1]. Juho Kellosoalo, et al. Crystallization and preliminary X-ray analysis of membrane- bound pyrophosphatases. Mol Membr Biol. 2013 Feb;30(1):64-74.
- [2]. Muhammad Ehsan, et al. New Malonate-Derived Tetraglucoside Detergents for Membrane Protein Stability. ACS Chem Biol. 2020 Jun 19;15(6):1697-1707.

For reference only
For Research Use Only. Not for Diagnostic or Therapeutic Use.