

Slo2.2 Antibody

Slo2.2 Antibody, Clone S3-26 Catalog # ASM10199

Specification

Slo2.2 Antibody - Product Information

Application WB, IHC
Primary Accession O9Z258
Other Accession NP_068625
Host Mouse
Isotype IgG1

Reactivity Human, Mouse, Rat

Clonality Monoclonal

Description

Mouse Anti-Rat Slo2.2 Monoclonal IgG1

Target/Specificity

Detects ~140kDa. Weak human detection.

Other Names

KCNT1 Antibody, potassium channel subfamily T member 1 Antibody, bA100C15.2 Antibody, SLACK Antibody, KCa4.1KIAA1422FLJ41282 Antibody

Immunogen

Fusion protein amino acids 1168-1237 of rat Slo2.2 (Slack)

PurificationProtein G Purified

Storage -20°C

Storage Buffer

PBS pH7.4, 50% glycerol, 0.09% sodium azide

Shipping Temperature Blue Ice or 4°C

Certificate of Analysis

 $1 \mu g/ml$ of SMC-323 was sufficient for detection of Slo2.2 in $10 \mu g$ of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.

Cellular Localization

Cell Membrane

Slo2.2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry




- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Slo2.2 Antibody - Images




Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Slo2.2 Potassium Channel Monoclonal Antibody, Clone S3-26 (ASM10199). Tissue: HaCaT cells. Species: Human. Fixation: Cold 100% methanol for 10 minutes at -20°C. Primary Antibody: Mouse Anti-Slo2.2 Potassium Channel Monoclonal Antibody (ASM10199) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Dotty staining around nucleus and some in cytoplasm.



Western Blot analysis of Rat brain membrane lysate showing detection of Slo2.2 Potassium Channel protein using Mouse Anti-Slo2.2 Potassium Channel Monoclonal Antibody, Clone S3-26 (ASM10199). Load: 15 μ g. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-Slo2.2 Potassium Channel Monoclonal Antibody (ASM10199) at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.





Immunohistochemistry analysis using Mouse Anti-Slo2.2 Potassium Channel Monoclonal Antibody, Clone S3-26 (ASM10199). Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-Slo2.2 Potassium Channel Monoclonal Antibody (ASM10199) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Suprabasal epidermal staining. Hair follicles negative.

Slo2.2 Antibody - Background

Slo2.2 is a novel member of the mammalian Slo potassium channel gene family. Slo2 channels may contribute to the resting potentials of cells that control their basal level of excitability (1). They also have sensors that couple channel activity to the intracellular concentrations of Na+ and Cl- (2).

Slo2.2 Antibody - References

- 1. Santi C.M., et al. (2006) J Neuroscience, 26(19): 5059-5068.
- 2. Yuan A., et al. (2003) Neuron 37: 765-773.