

ENaC alpha Antibody

ENaC alpha Antibody, Clone 2G4 Catalog # ASM10163

#### Specification

# **ENaC alpha Antibody - Product Information**

Application	IHC, WB
Primary Accession	<u>Q6IRJ1</u>
Other Accession	<u>NP_113736</u>
Host	Mouse
Isotype	lgG2b
Reactivity	Mouse, Rat
Clonality	Monoclonal
Format	APC
Description	
Mouse Anti-Rat ENaC alpha Monoclonal IgG2b	

**Target/Specificity** Detects ~85kDa.

#### **Other Names**

SCNN1A Antibody, Epithelial Sodium Channel-α Antibody, Epithelial Sodium Channel alpha Antibody, Alpha ENaC 2 Antibody, Alpha ENaC Antibody, Alpha NaCH Antibody, Alpha-ENaC Antibody, Amiloride sensitive epithelial sodium channel alpha subunit Antibody, Amiloride sensitive sodium channel subunit alpha Antibody, Amiloride-sensitive sodium channel subunit alpha Antibody, ENaCa Antibody, ENaCalpha Antibody, Epithelial Na(+) channel subunit alpha Antibody, Epithelial Na+ channel subunit alpha Antibody, FLJ21883 Antibody, Nonvoltage gated sodium channel 1 subunit alpha Antibody, Nonvoltage-gated sodium channel 1 subunit alpha Antibody, SCNEA Antibody, SCNN 1 Antibody, SCNN1 Antibody, SCNN1A Antibody, SCNNA\_HUMAN Antibody, Sodium channel nonvoltage gated 1 alpha Antibody

Immunogen

Synthetic peptide from the N-terminal of Rat ENaC alpha (aa. 46-68)

Purification Protein G Purified

Storage Storage Buffer PBS pH7.4, 50% glycerol, 0.09% sodium azide -20ºC

Blue Ice or 4ºC

Shipping Temperature

**Certificate of Analysis** A 1:1000 dilution of SMC-239 was sufficient for detection of ENaC alpha in 15  $\mu$ g of Mouse whole kidney lysate by ECL immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.

**Cellular Localization** Apical Cell Membrane




# ENaC alpha Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

#### ENaC alpha Antibody - Images




Immunohistochemistry analysis using Mouse Anti-ENaC alpha Monoclonal Antibody, Clone 2G4 (ASM10163). Tissue: Kidney. Species: Rat. Fixation: Paraffin-embedded formalin-fixed. Primary Antibody: Mouse Anti-ENaC alpha Monoclonal Antibody (ASM10163) at 1:100. Secondary Antibody: Goat Anti-Mouse ATTO 488 (green). Localization: Intercalated cells. Aquaporin 2 Antibody staining in red.



Immunohistochemistry analysis using Mouse Anti-ENaC alpha Monoclonal Antibody, Clone 2G4 (ASM10163). Tissue: Kidney (cortex). Species: Mouse. Primary Antibody: Mouse Anti-ENaC alpha Monoclonal Antibody (ASM10163) at 1:150. Localization: Collecting duct principal cells. Magnification: 60X.





Western Blot analysis of Mouse Whole kidney homogenates showing detection of ~85kDa ENaC alpha protein using Mouse Anti-ENaC alpha Monoclonal Antibody, Clone 2G4 (ASM10163). Lane 1: Molecular Weight Ladder (MW). Lane 2: Low-salt diet. Lane 3: Normal-salt diet. Load: 20 µg. Primary Antibody: Mouse Anti-ENaC alpha Monoclonal Antibody (ASM10163) at 1:1000. Predicted/Observed Size: ~85kDa.

# ENaC alpha Antibody - Background

The Epithelial Sodium Channel (ENaC) is a membrane ion channel permeable to Na+ ions. It is located in the apical plasma membrane of epithelia in the kidneys, lung, colon, and other tissues where it plays a role in trans epithelial Na+-ion transport (1). Specifically Na+ transport via ENaC occurs across many epithelial surfaces, and plays a key role in regulating salt and water absorption (2).

ENaCs are composed of three structurally related subunits that form a tetrameric channel, alpha, beta, and gamma. The expression of its alpha and beta subunits is enhanced as keratinocytes differentiate (3, 4). The beta and gamma-ENaC subunits are essential for edema fluid to exert its maximal effect on net fluid absorption by distal lung epithelia(5). And it has been concluded that the subunits are differentially expressed in the retina of mice with ocular hypertension, therefore the up-regulation of alpha-ENaC proteins could serve as a protection mechanism against elevated intraocular pressure (6).

# ENaC alpha Antibody - References

- 1. Kakizoe Y., et al. (2009) J Hpyertens. 27(8): 1679-1689.
- 2. Gu Y. (2008) J Cell Physiol. 216(2):453-457.
- 3. Bruns J.B. (2003) Am J Physiol Renal Physiol. 285(4): F600-F609.
- 4. Mauro T., et al. (2002) J Invest Dermatol. 118(4): 589-594.
- 5. Elias N., et al. (2007) Am J Physiol Lung Cell Mol Physiol. 293(3): L537-45.
- 6. Dyka F.M., May C.A. and Enz R. (2005) J Neurochem. 94(1): 120-128.