

Notch1 Antibody
Notch1 Antibody, Clone S253-32
Catalog # ASM10264**Specification**

Notch1 Antibody - Product Information

Application	WB
Primary Accession	Q01705
Other Accession	NP_032740.3
Host	Mouse
Isotype	IgG1
Reactivity	Mouse, Rat
Clonality	Monoclonal
Format	PerCP

Description

Mouse Anti-Mouse Notch1 Monoclonal IgG1

Target/Specificity

Detects >270kDa, ~120kDa and small fragments due to proteolysis. Does not cross-react with Notch 2 or Notch3.

Other Names

Neurogenic locus notch homolog protein 1 Antibody, hN1 Antibody, Translocation associated notch protein Antibody, TAN1 Antibody, TAN-1 Antibody, Motch A Antibody, mT14 Antibody, p300 Antibody, Motch Antibody, Mis6 Antibody, Neurogenic locus notch protein homolog Antibody, NICD Antibody, Notch 1 intracellular domain Antibody, Notch gene homolog 1 (Drosophila) Antibody, Notch homolog 1 Antibody, translocation-associated (Drosophila) Antibody, NOTCH Drosophila Antibody, homolog of 1 Antibody, Translocation associated notch homolog Antibody, Translocation associated notch protein TAN 1 Antibody, xotch Antibody

Immunogen

Fusion protein amino acids 20-216 (extracellular N-terminus, EGF-like domains 1-5) of mouse Notch1

Purification

Protein G Purified

Storage **-20°C****Storage Buffer**

PBS pH7.4, 50% glycerol, 0.09% sodium azide

Shipping Temperature

Blue Ice or 4°C**Certificate of Analysis**

1 µg/ml of SMC-430 was sufficient for detection of Notch1 in 20 µg of rat brain membrane lysate and assayed by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.

Cellular Localization


Cell Membrane | Nucleus

Notch1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Notch1 Antibody - Images



Western Blot analysis of Rat Brain Membrane showing detection of ~270 kDa Notch1 protein using Mouse Anti-Notch1 Monoclonal Antibody, Clone N253/32 (ASM10264). Lane 1: MW Ladder. Lane 2: Rat Brain Membrane (10 µg). . Load: 10 µg. Block: 5% milk. Primary Antibody: Mouse Anti-Notch1 Monoclonal Antibody (ASM10264) at 1:1000 for 1 hour at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:200 for 1 hour at RT. Color Development: TMB solution for 10 min at RT. Predicted/Observed Size: ~270 kDa.

Notch1 Antibody - Background

Notch is synthesized in the endoplasmic reticulum as an inactive form which is proteolytically cleaved by a furin-like convertase (S1 cleavage) in the trans-golgi network before it reaches the plasma membrane to yield an active, ligand-accessible form. Cleavage results in a C-terminal fragment N(TM) and a N-terminal fragment N(EC). Following ligand binding, it is cleaved (S2 cleavage) by TNF-alpha converting enzyme (TACE) to yield a membrane-associated intermediate fragment called Notch extracellular truncation (NEXT). This fragment is then cleaved by presenilin-dependent gamma-secretase (S3 cleavage) to release the intracellular domain (NICD) from the membrane.

Notch1 Antibody - References

1. Swiatek P.J., et al. (1994) Genes Dev. 8: 707-719.
2. Simpson P. (1994) The Notch Receptors. Austin, TX: R.G. Landes Company.
3. Lindsell C.E., et al. (1995) Cell 80: 909-917.

4. Uyttendaele H., et al. (1996) *Development* 122: 2251-2259.
5. Girard L., et al. (1996) *Genes Dev.* 10: 1930-1944.